

2023

NSARB-2023-001

Nova Scotia Aquaculture Review Board

IN THE MATTER OF: Applications made by **KELLY COVE SALMON LTD.** for a **BOUNDARY AMENDMENT** and **TWO NEW MARINE FINFISH AQUACULTURE LICENSES** and **LEASES** for the cultivation of **ATLANTIC SALMON (Salmo salar)** - **AQ#1205x, AQ#1432, AQ#1433** in **LIVERPOOL BAY, QUEENS COUNTY.**

Kelly Cove Salmon Ltd.

APPLICANT

-and-

Minister of Fisheries and Aquaculture

PARTY

-and-

Kwilmu'kw Maw-Klusuaqn Negotiation Office (KMKNO)

FIRST INTERVENOR

-and-

Queens Recreational Boating Association (Brooklyn Marina)

SECOND INTERVENOR

22 Fishermen of Liverpool Bay

THIRD INTERVENOR

Region of Queens Municipality

FOURTH INTERVENOR

Protect Liverpool Bay Association


FIFTH INTERVENOR

Affidavit of Dr. Chris McKindsey

I, Christopher W. McKindsey, of Rimouski, Quebec, affirm and give evidence as follows:

1. I am the Research Scientist and Section Head, Aquaculture and Aquatic Invasive Species Section in the Department of Fisheries and Oceans Canada (DFO). I attach a copy of my Resume hereto as **Exhibit "A"**.
2. I have personal knowledge of the evidence affirmed in this Affidavit except where otherwise stated to be based on information and belief.
3. I state, in this Affidavit, the source of any information that is not based on my own personal knowledge, and I state my belief of the source.
4. I have been asked to describe my recent research project respecting lobster telemetry in Liverpool Bay, Nova Scotia.
5. My project partners and I have co-authored a manuscript presenting the preliminary results of my research on lobster telemetry in Liverpool Bay. The results of this research are referred to as "preliminary" as this material has not yet been published and peer reviewed in academic scientific literature. This manuscript is attached hereto as **Exhibit "B"**.
6. I was not physically present before Mrs. Campbell when I affirmed this affidavit. I was linked with Mrs. Campbell using video conferencing technology.

Sworn/Affirmed before me by
videoconference from Rimouski, Quebec,
(location of affiant) to Halifax, Nova Scotia
(location of lawyer taking oath) on the 22
day of January 2024.


Alison W. Campbell
A Barrister of the Supreme Court
of Nova Scotia

McKindsey,
Chris


Digitally signed by McKindsey, Chris
DN: C=CA, O=GC, OU=DFO-MPO, CN="McKindsey, Chris"
Reason: I approve this document
Location: Rimouski, Qc
Date: 2024.01.22 13:08:13-05'00'
Foxit PDF Editor Version: 13.0.1

Christopher W. McKindsey

2023

NSARB-2023-001

This is Exhibit "A" referred to in the
Affidavit of Christopher McKindsey
affirmed before me by videoconference
on January 22, 2024

A large black rectangular redaction covers the signature area. A small blue mark is visible below the redaction.

Signature

ALISON W. CAMPBELL

A Barrister of the Supreme Court of Nova Scotia

Christopher W. McKindsey

Research Scientist and Section Head, Aquaculture and Aquatic Invasive Species Section
Fisheries and Oceans Canada, Maurice-Lamontagne Institute
PO Box 1000, 850 route de la Mer, Mont-Joli, QC, G5H 3Z4, Canada

[REDACTED]@dfo-mpo.gc.ca

EDUCATION: Postdoctoral fellow (Marine Ecology, University of Sydney, 1999-2001)
Ph.D. (Marine Ecology, Université Laval, 1999)
M.Sc. (Parasitology, Concordia University, 1993)
B.Sc. (Ecology/Zoology, Concordia University, 1990)

OVERVIEW

- ~ Thirty years' experience in aquatic ecology, including research on rocky intertidal and pelagic marine systems, fresh water wetlands, restoration ecology and environmental impact studies, aquaculture-environment interactions, and exotic species;
- ~ extensive experience in managing research projects and research teams in terms of setting research goals, human resources management, equipment procurement and management, and financial management;
- ~ solid communications skills for various audiences (scientific communications: peer-reviewed articles/presentations, communication with clients, and regular contributor to information sources for non-scientists – interviews, newsletters, public speaking, etc.);
- ~ proven ability to secure funding through competitive grants (Fisheries and Oceans Canada, NSERC, industry);
- ~ international, national, and local experience and collaborations, with ministerial, academic, First Nations, and industry partners;
- ~ adjunct professor at ISMER and Université Laval, and co-director of students at multiple universities.

RESEARCH INTERESTS

- ~ Current research examines the influence of humans on the marine environment with an emphasis on four related areas:
 1. Aquaculture-Environment Interactions. Current work examines impact of bivalves on the pelagic ecosystem, interactions between bivalve and salmon aquaculture and CRA species, such as lobster and crab using modelling approaches.
 2. Invasive species. Current work examines the importance of shipping / boating as vectors for the establishment of invasive species and on the physiology of invasive species, including *Mya arenaria* and *Littorina littorea*, and species distribution modeling.
 3. Acoustic telemetry. Work focuses on the movement of decapods, other macroinvertebrates, and fish, often in relation to aquaculture sites (fish and decapods) but also with respect to habitat structures (decapods, whelks, echinoderms).
 4. Other. Much work focuses on benthic ecology in the Canadian Arctic in support of AIS and baseline biodiversity monitoring, including mapping benthic ecological assemblages and interactions between habitats and fish / invertebrates.

RESEARCH MANAGEMENT

- ~ Section Head for Coastal and Benthic Ecology Section (2010-2014, 2018-today; 10 permanent and 1-10 temporary employees):
 1. Oversee the research programme for the section to ensure it meets client needs, including setting research priorities following discussions with stakeholders, seeking internal/external funds through grants, participation in national and other committees.
 2. Managed personnel within section, including contributing to developing evaluation tools for hiring Research Scientists (RES) across Canada and specific criteria to hire a new RES and Biologist and many temporary staff, dividing tasks/needs within section, managing many graduate/summer students, and ensuring partnerships are fostered with internal and external expertise.
 3. Oversaw finances for multiple budgets (\$300K-\$1200K yr⁻¹ for research expenses plus staffing expenses for the past 10 years) both within and outside of the Department (e.g., DFO, industry, and NSERC funds – all competitive research funds).
 4. Ensured that all required training (e.g., first-aid, heavy vehicle, finances) was up to date for all permanent and temporary staff and students. Encouraged team to follow appropriate advanced training, sought it out and secured resources to pay for it.

INFLUENCE

Aquaculture

1. Invited to submit multiple projects to ACRDP, PARR, and Galway programmes within DFO as well as through NSERC (AquaNet, CAISN, CHONe) and with industry (other than ACRDP; BC Salmon Farmers Association);
2. Chair (2008-2012) of ICES Working Group on Environmental Interactions of Mariculture (WGEIM), past member of Working Group on Aquaculture (WGAQUA), member of WGECCA (Ecological Carrying Capacity for Aquaculture) and WGREIA (Risk assessment of Environmental Interactions of Aquaculture);
3. Invited to contribute to a book on goods and services of marine bivalves (emphasis on farmed and natural populations) (Springer);
4. Invited to contribute to 2 chapters for Encyclopedia of Sustainability Science and Technology (Springer);
5. Numerous collaborative projects locally, nationally, and internationally.

Invasive Species

1. Theme lead for NSERC Canadian Healthy Oceans Network CHONe (Ecosystem function) and CHONe II (Cumulative stressors);
2. Eastern node lead for NSERC Canadian Aquatic Invasive Species Network (CAISN I and CAISN II);
3. Numerous collaborative projects locally, nationally, and internationally;

4. Member of ICES Working Groups on Aquatic Invasive Species, including the Ballast and Other Ship Vectors (WGBOSV) and Introductions and Transfers of Marine Organisms (WGITMO).

RECENT PRIMARY PUBLICATIONS;

- English G, Wilson BM, Lawrence MJ, Hawkes JP, Hardie DC, Daniels J, Carr J, Rycroft C, Whoriskey FH, **McKindsey CW**, Trudel M (accepted) Determining early marine survival and marine mammal predation on acoustically-tagged Atlantic salmon (*Salmo salar*) post-smolts. *Can J Fish Aquat Sci*
- Goldsmith J, **McKindsey CW**, Schlegel RW, Deslauriers D, Howland KL (accepted) Predicting distribution overlap of interacting species: Match or mismatch in a warmer and invaded Canadian Arctic? *Elementa: Science of the Anthropocene*
- Nadalini JB, Lees KJ, Lavoie M-F, MacGregor KA, **McKindsey CW** (accepted) An evaluation of acoustic telemetry as a method to study the movements of Asteroidea (*Asterias rubens*). *Anim Biotelemetry*
- Dreuou E, Beauchesne D, Daigle RM, Carrière J, Noisette F, **McKindsey CW**, Archambault P (2023) Multiple human activities in coastal benthic ecosystems: introducing a metric of cumulative exposure. *Elementa: Science of the Anthropocene* 11:1
- Finnis S, Guyondet T, **McKindsey CW**, Filgueira R, Areseneau J, Barrell J, Gibb O, Gallardi D, Milne R, Goodwin C, Macdonald TA, Duhaime J, Lacoursière-Roussel A (2023) Guidance on sampling effort to monitor mesozooplankton communities at bivalve aquaculture sites using an optical imaging system. *Can Tech Rep Fish Aquat Sci* 3581:vii + 98
- Fisher J, Angel D, Callier M, Cheney D, Filgueira R, Hudson B, **McKindsey CW**, Milke L, Moore H, O'Beirn F, O'Carroll J, Rabe B, Telfer T, Byron CJ (2023) Ecological carrying capacity in mariculture: Consideration and application in geographic strategies and policy. *Mar Policy* 150:105516
- Gianasi BL, **McKindsey CW**, Tremblay R, Comeau LA, Drolet D (2023) Plankton depletion by mussel grazing negatively impacts the fitness of lobster larvae. *Aquaculture* 574:739659
- Goldsmith J, Clark HA, **McKindsey CW**, Stewart DB, Howland KL (2023) Screening for high-risk marine invaders in the Hudson Bay Region, Canadian Arctic: Compilation of background information, rationale, and references used to answer questions with the Canadian Marine Invasive Species Tool (CMIST). *Can Data Rep Fish Aquat Sci* 1373:vi + 344
- Kingsbury MV, Hamoutene D, Kraska P, Lacoursière-Roussel A, Page F, Coyle T, Sutherland T, Gibb O, **McKindsey CW**, Hartog F, Neil S, Chernoff K, Wong D, Law BA, Brager L, Baillie SM, Black M, Bungay T, Gaspard D, Hua K, Parsons GJ (2023) Investigation on the association of anti-leech drugs and antibiotics used during finfish production with organic enrichment in marine sediments at Canadian aquaculture sites. *Mar Pollut Bull* 188:114654
- Lawrence MJ, Wilson BM, Reid GK, Hawthorn C, English G, Black M, Leadbeater S, **McKindsey CW**, Trudel M (2023) The fate of intracoelomic acoustic transmitters in Atlantic salmon (*Salmo salar*) post-smolts and wider considerations for causal factors driving tag retention and mortality in fishes. *Anim Biotelemetry* 11:40
- Lees KJ, Lavoie M-F, MacGregor KA, Simard É, Drouin A, Comeau LA, **McKindsey CW** (2023) Movement of American lobster *Homarus americanus* and rock crabs *Cancer irroratus* around mussel farms in Malpeque Bay, Prince Edward Island, Canada. *Aquacult Environ Interact* 15:179–193
- MacGregor KA, Lavoie M-F, Robinson SMC, Simard É, **McKindsey CW** (2023) Lab and field evaluation of tagging methods for the use of acoustic telemetry to observe sea urchin movement behaviour at ecologically relevant spatio-temporal scales. *Anim Biotelemetry* 11:3
- Mignucci A, Forget F, Villeneuve R, Derridj O, **McKindsey CW**, McKenzie D, Bourjea J (2023) Residency, home range and inter-annual fidelity of three coastal fish species in a Mediterranean coastal lagoon. *Estuar Coast Shelf Sci* 292:108450
- Schull Q, Viblanç V, Metral L, Leclerc L, Romero D, Pernet F, Quézé C, Derolez V, Munaron D, **McKindsey CW**, Saraux C, Bourjea J (2023) An integrative perspective on fish health: environmental and anthropogenic pathways affecting fish stress. *Mar Pollut Bull* 194:115318
- Bringloe TT, Wilkinson DP, Goldsmith J, Savoie AM, Filbee-Dexter K, Macgregor KA, Howland KL, **McKindsey CW**, Verbrugge H (2022) Arctic marine forest distribution models showcase potentially severe habitat losses for cryophilic species under climate change. *Glob Change Biol* 28:3711-3727
- Dhifallah F, Rochon A, Simard N, **McKindsey CW**, Gosselin M, Howland KL (2022) Dinoflagellate communities in high-risk Canadian Arctic ports. *Estuar Coast Shelf Sci* 266:107731
- Drolet D, Riley C, Robert S, Estrada R, Gianasi BL, **McKindsey CW** (2022) Effect of aquaculture-related diets on the long-term performance and condition of the rock crab, *Cancer irroratus*. *Front Mar Sci* 9:865390
- Ferrario F, Araújo CAS, Bélanger S, Bourgault D, Carrière J, Carrier-Belleau C, Dreuou E, Johnson LE, Juniper K, Mabit R, **McKindsey CW**, Ogston L, Picard MMM, Saint-Louis R, Saulnier-Talbot É, Shaw J-L, Templeman N, Therriault TW, Tremblay J-E, Archambault P (2022) Holistic environmental monitoring in ports as an opportunity to advance sustainable development, marine science and social inclusiveness. *Elementa: Sci Anthropocene* 10:1
- Filbee-Dexter K, MacGregor KA, Lavoie C, Garrido I, Goldsmith J, Castro de la Guardia L, Howland KL, Johnson LE, Konar B, **McKindsey CW**, Mundy CJ, Schlegel RW, Archambault P (2022) Sea ice and substratum shape extensive kelp forests in the Canadian Arctic. *Front Mar Sci* 9:754074
- Gianasi BL, Goldsmith J, Archambault P, **McKindsey CW**, Holovachov O, Howland KL (2022) Biodiversity of macrobenthic nematodes in the intertidal and shallow subtidal zones in the eastern Canadian Arctic. *Polar Biol* 45:225-242
- Gianasi BL, **McKindsey CW**, Howland KL (2022) Coastal biodiversity of epibenthic assemblages in eastern Canadian Arctic: Baseline mapping for management and conservation. *Front Mar Sci* 9:873608
- Lavoie M-F, Simard É, Drouin A, Archambault P, Comeau LA, **McKindsey CW** (2022) Movements of American lobster (*Homarus americanus*) associated with offshore mussel (*Mytilus edulis*) aquaculture. *Aquacult Environ Interact* 14:189-204
- Ma KCK, **McKindsey CW**, Johnson LE (2022) Detecting rare species with passive sampling tools: optimizing the duration and frequency of sampling for benthic taxa. *Front Mar Sci* 9:809327
- Quinn BK, Trudel M, Wilson BM, Carr J, Daniels J, Haigh S, Hardie DC, **McKindsey CW**, O'Flaherty-Sproul M, Simard É, Page F (2022) Modelling the effects of currents and migratory behaviours on the dispersal of Atlantic salmon (*Salmo salar*) post-smolts in a coastal embayment. *Can J Fish Aquat Sci* 79:2087–2111
- Riley C, Drolet D, Goldsmith J, Hill JM, Howland KL, Lavoie M-F, McKenzie CH, Simard N, **McKindsey CW** (2022) Experimental analysis of survival and recovery of ship fouling mussels during transit between marine and freshwaters. *Front Mar Sci* 8:808007
- Robichaud L, Archambault P, Desrosiers G, **McKindsey CW** (2022) Influence of suspended mussel aquaculture and an associated invasive ascidian on benthic macroinvertebrate communities. *Water* 14:2751
- Sean A-S, Drouin A, Archambault P, **McKindsey CW** (2022) Influence of an offshore mussel aquaculture site on the distribution of epibenthic macrofauna in Îles de la Madeleine, eastern Canada. *Front Mar Sci* 9:859816

2023

NSARB-2023-001

This is Exhibit "B" referred to in the
Affidavit of Christopher McKindsey
affirmed before me by videoconference
on January 22, 2024

A large black rectangular redaction box covering the signature of the witness.

Signature

ALISON W. CAMPBELL

A Barrister of the Supreme Court of Nova Scotia

1 **Impact of finfish farms in eastern Canada (Nova Scotia) on American lobster and rock crab**
2 **distribution**

3 *Running page head: Lobster and crab movements around finfish farms*
4

5 Marie-France Lavoie¹, Nathaniel Feindel², Ryan A. Horricks³, Shawn M. C. Robinson⁴, and
6 Christopher W. McKindsey^{1*}

7 ¹ Fisheries and Oceans Canada, Maurice Lamontagne Institute, Mont-Joli, Québec, G5H 3Z4, Canada

8 ² Nova Scotia Department of Fisheries and Aquaculture, Shelburne, Nova Scotia, B0T 1W0, Canada

9 ³ Centre for Marine Applied Research, 27 Parker Street, Dartmouth, Nova Scotia, B2Y 4T5, Canada

10 ⁴ Fisheries and Oceans Canada, Saint Andrews Biological Station, Saint Andrews, New Brunswick, E5B 2L9,
11 Canada

12 * Corresponding author: chris.mckindsey@dfo-mpo.gc.ca
13

14 **ABSTRACT**

15

16 **Keywords:** American lobster; Rock crab; Abundance; Acoustic telemetry

17

18

19

20 1. INTRODUCTION

21 Aquaculture is increasing worldwide and in Canada, particularly regarding finfish culture
22 (Fisheries and Oceans Canada 2021, FAO 2022). Finfish farms are known to attract wild animals
23 and interactions between fish farms and wild fishes have been well documented (see Callier *et al.*
24 2018; Barrett *et al.* 2019). Many mechanisms have been identified that attract and repel wild fish to
25 and from fish farms: waste feed, farms acting as fish aggregation devices (FADs) or artificial reefs
26 (ARs), biofouling communities on the nets, secondary attraction of predators, benthic changes, and
27 husbandry practices (Callier *et al.* 2018; Barrett *et al.* 2019). Less is known about how fish farms
28 attract or repel invertebrates, despite the importance of several commercial invertebrate species,
29 including the American lobster (*Homarus americauns*) (Florko *et al.* 2021).

30 American lobster is the most valuable fishery in eastern Canada (ca. \$CAN 2.05 B in 2021;
31 Fisheries and Oceans Canada 2021). There are increasing concerns about the influence of finfish
32 aquaculture sites on lobster distribution patterns and potential changes in food sources that may
33 impact their condition, biology, and catchability (reviewed in Horricks *et al.* 2022). The perception
34 is often that fish farms may negatively impact lobster abundance and condition (Wiber *et al.* 2012;
35 Loucks *et al.* 2014; Grant *et al.* 2016; Loucks *et al.* 2016; Milewski *et al.* 2018). In addition, the
36 rock crab is also fished commercially in eastern Canada, and although the landings of this species
37 are dwarfed by the landed value of lobster, it is of importance in several areas (Fisheries and Oceans
38 Canada 2022). The preferred prey of lobster is rock crab (Gendron *et al.* 2001; Hanson 2009) and
39 unpublished work from southwest New Brunswick suggests that there are clear differences in
40 associations of rock crab and lobster to salmon farm sites making a comparison between the two
41 species of interest.

42 According to Loucks *et al.* (2014), fish farm activities may impact crustaceans close to
43 salmonid farms by altering their movement and behaviour. Work on lobster movement near mussel
44 farms in Iles-de-la-Madeleine and Prince Edward Island (Lavoie *et al.* 2022; Lees *et al.* 2023). In
45 Iles-de-la-Madeleine, lobsters had little affinity to mussel farms, neither setting up territories within
46 farm limits nor remaining in them for extended periods. Work in Prince Edward Island showed that
47 lobsters may use mussel farms for foraging and shelter and that they entered and exited farms
48 frequently, sometimes multiple times a day (Lees *et al.* 2023). In contrast, rock crabs mostly
49 remained within mussel farm lease boundaries, staying mostly directly below culture structures
50 (Lees *et al.* 2023).


51 This study evaluated the abundance and movement of decapods in the near vicinity of two
52 salmonid aquaculture leases in Port Mouton and Liverpool Bay, Nova Scotia, Canada, using direct
53 observation of animals and acoustic telemetry. The study was done over a full production cycle in
54 Liverpool Bay, starting with a fallow year (no fish on site) and subsequently evaluating decapod
55 interactions with a farm site at different production stages (fallow, 1-year old fish and 2-year old
56 fish prior to harvesting). Previous work done around the Port Mouton Atlantic salmon/rainbow trout
57 farm suggested it impacts the distribution of market and berried lobster (Loucks *et al.* 2014;
58 Milewski *et al.* 2018), making it of interest in the present study. Results from this study will inform
59 managers on the spatial interactions between two economically important activities in eastern
60 Canada: marine farming of salmonids and lobster/crab fisheries. It is hoped that results from this
61 study will foster the development of a sustainable salmonid aquaculture industry while allowing
62 the continued use of fisheries resources by providing an evidence-based understanding of the links
63 between these two activities.

64

65 2. MATERIALS AND METHODS

66 Study areas

67 The study was conducted in Port Mouton Bay (43°54'15"N, 64°47'31"N) and in Liverpool
68 Bay (44°01'59"N, 64°39'31"W), Nova Scotia (NS), Canada in 2019, 2020, and 2021 (Figure 1).
69 Port Mouton Bay is partially sheltered with depth varying from 8 to 18 m with the benthic substrate
70 characterized by a mix of sand, gravelly sand, and mud (Piper *et al.* 1986). Liverpool Bay is 4.5 km
71 long, 2.6 km wide, and open to the ocean. Coffin Island protects the northeast entrance to the bay
72 and provides shelter for the current Atlantic salmon pen aquaculture site and proposed sites. Benthic
73 substrate in Liverpool is principally characterised by sand (in the middle of the bay), rock, and
74 gravel (McKee *et al.* 2021). Neither aquaculture site was in use in 2019. Since no further salmonid
75 aquaculture was planned for the Port Mouton site following 2019 (CWM personal communications
76 with the Nova Scotia Department of Fisheries and Aquaculture), the acoustic telemetry study was
77 not continued at this location. As of the writing of this manuscript, there is a proposal to expand
78 finfish aquaculture in Liverpool, NS (Figure 1). During the 2019 field season, the physical structure
79 of the cages (i.e. the floating Polarcirkel fish pens; Klepp stasjon, Norway) were on the Liverpool
80 site (7 × 2 cages, near Coffin Island, NS) and only some buoys and ropes were visible in the north
81 site (at the northwest of the Spectacle Islands) in Port Mouton, although no fish were in cages at
82 either site. In 2020, fish were added to the cages in Liverpool and in 2021 the cages contained 2-
83 year-old Atlantic salmon.



84

85 **Figure 1.** A) Location of the two study sites (black dots) in southern Nova Scotia (Canada). B)

86 Location of the Port Mouton receivers in 2019 where the dashed outline represents the previously

87 farmed site, and in Liverpool in C) 2019, D) 2020, and E) 2021. The grey dots represent the acoustic

88 receivers in the study sites. For C-E, the 1205x polygon represents the finfish lease and the 1432-

89 1433 polygons represent the proposed culture sites (1432 - Fralick Cove and 1433 - Mersey Point).

90

91 **Observational sampling**

92 In July 2019, a total of 35 transects (50 m long x 2 m wide) were swam by SCUBA divers
93 in five distinct areas (farm site, Fralick Cove, Mersey Point, and two reference sites) in Liverpool
94 Bay. Divers collected all decapods on encounter and brought them to the surface where they were
95 sexed, measured, and counted. In Port-Mouton, SCUBA divers collected decapods on encounter
96 along 50 transects in 5 distinct areas (Proposed farm site, 2 reference sites near the proposed farm,
97 and 2 reference sites to the southeast of these). No transects were done at any location in 2020 due
98 to travel restrictions caused by the COVID-19 pandemic making such voyages impossible. In 2021,
99 30 transects were sampled in Liverpool Bay (farm site, Fralick Cove, and Mersey Point) with all
100 transects being done in the same way as in 2019.

101 **Acoustic telemetry design and tag deployments**

102 Three receiver models were deployed during the study: VR2W, VR2Tx, and VR2AR (69
103 kHz, Innovasea). Receivers were deployed in a grid that satisfied the criteria for the VEMCO
104 Positioning System (VPS), fine-scale movement analysis (Orrell & Hussey 2022, Espinoza et al.
105 2011). At the beginning of the study, a range test was done prior the deployment of the acoustic
106 receiver grid to determine optimal receiver spacing. A synchronisation tag (model V13, 36 mm
107 long and 13 mm diameter, transmission delay between 500-700 s, Innovasea) was deployed 1 m
108 above each VR2W receiver; VR2Tx and VR2AR receivers contain synchronisation tags
109 (transmission delay between 540-660 s, Innovasea) within the units. Two or three reference tags
110 (V9, Innovasea) were used at each site and placed where most receivers could record the signal.

111

112

113 2019

114 Figure 1 (B and C) shows the design for each location bay with the position of acoustic
115 receivers. A substantial portion of Liverpool Bay was covered with receivers (Figure 1C) and three
116 areas (N = North – where the salmonid farm was, C = Center, and S = South) were studied in Port
117 Mouton (Figure 1B). This was a considerable sampling campaign with a total of 138 receivers
118 deployed (Liverpool = 81 and Port Mouton = 57) in 2019. Receivers were deployed between July
119 16 and 23 and recovered between November 19 and 22 2019, for a total of 124 days for Liverpool
120 and 121 days for Port Mouton. Each receiver was separated by 250 to 375 m from its neighbour.

121 A total of 47 rock crabs (34 M and 13 F) and 50 lobsters (29 M and 21 F) were captured and
122 tagged in Liverpool and 51 rock crabs (22 M and 29 F) and 50 lobsters (29 M and 21 F) captured
123 and tagged in Port Mouton (Table 1). All animals were measured (width for crabs and
124 cephalothorax length - CL - for lobsters) before being released at the same place they were caught
125 (farm site, Fralick Cove, and Mersey Point for Liverpool and north, center, and south sites for Port
126 Mouton). Animals were caught using commercial lobster traps or by SCUBA divers on encounter
127 and fitted on a boat with acoustic transmitters (Innovasea V9, 26 mm long and 9 mm diameter,
128 transmission delay between 180-300 s). Transmitters were attached to carapaces cleaned with
129 sandpaper followed by ethanol swabs) using ethyl 2-cyanoacrylate glue (LePage® Ultra Gel
130 Control® Super Glue) and released back into the study arrays. Transmitter attachment required
131 approximately 3 min for each animal, which were then released within about 10 min of being
132 brought to the surface.

133 2020

134 A total of 106 receivers was deployed in Liverpool Bay on June 29 to July 13 (Figure 1C)
 135 and were recovered November 3 to 4, 2020. Ten of these receivers were installed outside the bay
 136 at the northeast and deployed in two lines (Figure 1D). Each receiver was separated by 250 to 375
 137 m from its neighbour. One hundred animals were captured using commercial lobster traps, sexed,
 138 measured, and tagged (Table 1): 50 lobsters (30 M and 20 F) and 50 rock crabs (42 M and 8 F).

139 2021

140 Following the same methods used in 2019 and 2020, 104 receivers were deployed in
 141 Liverpool Bay on July 12 to 15 and recovered on November 9 to 10, 2021 (Figure 1E). A total of
 142 50 lobsters (27 M, 23 F) and 50 crabs (26 M, 24 F) was captured using commercial lobster traps
 143 and on encounter by SCUBA divers (Table 1).

144 **Table 1.** Number and sex of American lobster (*Homarus americanus*) and rock crab (*Cancer*
 145 *iroratus*) tagged each year and at each site in Nova Scotia, Canada.

Year	Species	Sex	Liverpool			Port Mouton		
			Farm	Fralick Cove	Mersey Point	North	Center	South
2019	Lobster	M	13	5	11	10	7	12
		F	12	0	9	10	8	3
	Rock crab	M	9	0	25	12	2	8
		F	11	0	2	9	13	7
2020	Lobster	M	18	5	7	-	-	-
		F	11	2	7	-	-	-
	Rock crab	M	36	4	2	-	-	-
		F	8	0	0	-	-	-
2021	Lobster	M	10	7	10	-	-	-
		F	10	8	5	-	-	-
	Rock crab	M	8	14	4	-	-	-
		F	12	1	11	-	-	-

146

147

148 **Data analysis**

149 Data were analyzed using the open-source statistical software R version 4.3.2 (R Core Team,
150 2023) and PRIMER-e (v. 7.0.21).

151 Acoustic telemetry data pre-processing

152 Prior to analysis, data were filtered using a linear regression to remove observations with
153 high positioning error (HPE) (Skerritt *et al.* 2015, Lees *et al.* 2020, Lavoie *et al.* 2022). HPE is a
154 relative measure of error and a calculated position with high HPE provides less precise information
155 on the position of an animal compared to a position with a lower HPE (Lees *et al.* 2023). A
156 regression was done for each site based on the synchronisation tag deployed at each site within each
157 year. HPE filtration was divided for each year of the study in Liverpool Bay. For 2019 in Liverpool
158 Bay, less than 20% of the synchronization tag data were lost by filtering by $HPE \leq 35$ ($r^2 = 0.99$).
159 The animal detections from 2019 were thus filtered with an $HPE \leq 35$ and a mean (\pm SE) position
160 error for individuals of 4.9 ± 0.01 m. For 2020, less than 1% of the synchronization tag data were
161 lost by filtering by $HPE \leq 25$ ($r^2 = 0.99$). Animal detections from 2020 were thus filtered with an
162 $HPE \leq 25$ giving a mean position error (\pm SE) of 12.4 ± 0.02 m. For the last year of the study in
163 Liverpool Bay, less than 5% of the sync tag data were lost by filtering by $HPE \leq 30$ ($r^2 = 0.88$). The
164 2021 animal detections were thus filtered with an $HPE \leq 30$ giving a mean position error (\pm SE) of
165 12.09 ± 0.01 m. Synchronization tag data from Port Mouton were filtered by $HPE \leq 30$ ($r^2 = 0.98$)
166 with a loss of less than 5% of the data. The mean position error (\pm SE) for Port Mouton animal
167 detections with $HPE \leq 30$ was 4.07 ± 0.008 m. The first 24 h of all tracking data were excluded at
168 all sites to minimize the impact of tagging on behavior (Lavoie *et al.* 2022).

169 Only animals with more than 200 detections over the entire deployment at each year were
170 kept for the analyses. A final filtration was used to remove data with individual walking speeds >
171 10 km d⁻¹ as these speeds are abnormal for lobster and rock crabs.

172 Movement parameters analyses

173 Animal track analyses were done using the adehabitatLT package (Calenge 2006). When
174 animals leave and then reenter the acoustic telemetry array, some gaps in the data may occur. To
175 avoid such gaps in residence time estimates, distance travelled, and walking speed trajectories were
176 split into separate bursts if the time between detections was >12 h and if the distance was >200 m
177 (Lees *et al.* 2023; Lavoie *et al.* 2022). Residence time and distance travelled correspond to the
178 cumulative time spent and cumulative distance travelled by each animal within the acoustic array,
179 respectively. Walking speed was estimated by dividing the distance interval from each step by the
180 time interval from the same step. Variations in residence time and the distance travelled for
181 Liverpool Bay were examined using 4-way ANOVAs with the fixed factors “Year” (3 levels: 2019,
182 2020, and 2021), tagging “Site” (3 levels: farm site, Fralick Cove, and Mersey Point), “Species” (2
183 levels: lobster and crab), and “Sex” (2 levels: M and F). Assumptions of homoscedasticity were
184 evaluated for ANOVA analyses using the Shapiro-Wilk test, as outlined in Quinn & Keough (2002).
185 Data were transformed, where necessary, to satisfy assumptions of ANOVA (square root for the
186 distance travelled). Variation in walking speed for Liverpool Bay was evaluated using
187 PERMANOVA (with 9999 permutations) as data transformations were unable to constrain the data
188 to meet the assumptions of ANOVA (Anderson 2001). The similarity matrix used to this end was
189 constructed based on Euclidean distances. The homogeneity of multivariate dispersion was
190 evaluated using PERMDISP and data transformed (square root). PERMANOVA analyses included
191 the same four factors as the ANOVA analyses.

192 Variation in movement parameters for Port Mouton individuals were evaluated using 3-way
193 ANOVAs with three fixed factors (“Site,” “Species,” and “Sex”). Assumptions of homoscedasticity
194 were evaluated for each ANOVA analysis using the Shapiro-Wilk test. Data were transformed to
195 satisfy assumptions of ANOVA (square root for the residence time and distance travelled, and log-
196 transformation for walking speed). Differences among treatment means of factors that were deemed
197 significant in the ANOVAs were evaluated using a posteriori Tukey multiple comparison tests.

198 Home range analyses

199 The 95% home range for the two species were calculated by kernel density estimations with
200 the “amt” package (Signer et al. 2019) in Liverpool Bay. With the home range results, an overlap
201 with the farm site was measured to determine the utilization percentage for each individual. To
202 compare the home range and the overlap results, PERMANOVA (based on 9999 permutations)
203 analyses were used to identify variation between four fixed factors (“Year,” “Site,” “Species,” and
204 “Sex”). Euclidian distances were used to construct the similarity matrix and data were transformed
205 as necessary (home range: square root; overlap: log+1). Differences among treatments were
206 determined using a posteriori pairwise comparisons, also using PERMANOVA. For Port Mouton,
207 home range data were analyzed with a 3-way ANOVA with three fixed factors (“Site,” “Species,”
208 and “Sex”) and with a fourth root transformation to satisfy assumptions. Differences among
209 treatment means of factors deemed significant in the ANOVA were evaluated using a posteriori
210 Tukey multiple comparison tests.

211 Animal abundance

212 Variation in animal abundance observed within transects in Liverpool Bay were examined
213 using PERMANOVA (with 9999 permutations) with the fixed factors “Year” (2 levels: 2019 and


214 2021), “Site” (3 levels: farm site, Fralick Cove, and Mersey Point), “Species” (3 levels: lobster,
215 rock crab, and Jonah crab), and “Sex” (3 levels: M, F, and juvenile) since data transformation were
216 unable to constrain the data to meet the assumptions of ANOVA (Anderson 2001). The similarity
217 matrix used was constructed based on Euclidean distances. The homogeneity of multivariate
218 dispersion was evaluated using PERMDISP and data transformed ($\log+1$). Differences among
219 treatments were determined using a posteriori pairwise comparisons, also using PERMANOVA.

220

221 3. RESULTS

222 Animal abundance

223 Three decapod species were observed in the transects at all sites: American lobster, rock
224 crab, and Jonah crab (*Cancer borealis*) (Figure 2). Abundance of Jonah crab was used for
225 abundance analysis, however, no Jonah crab were tagged with acoustic transmitters. Abundance
226 was found significantly different as a function of “Species,” “Sex,” and some factor interactions as
227 “Year \times Site \times Species” (Table 2). The number of lobsters observed decreased under fish farm from
228 2019 (fallow year) relative to 2021 (production year II) ($p = 0.0001$) whereas this decline was not
229 observed for rock crabs ($p = 0.4565$) (Figure 2; Table 2).



230

231 **Figure 2.** Abundance of American lobster (*Homarus americanus*), rock crab (*Cancer irroratus*),
 232 and Jonah crab (*Cancer borealis*) at each site in Liverpool Bay, Nova Scotia in 2019 and 2021.

233

234 **Table 2.** Results of PERMANOVAs for the animal abundance by year, tagging site, species, and sex.

235 Significant differences at $p < 0.05$ are highlighted in bold.

Source	Abundance			
	df	MS	F	p
Year	1	0.0071	0.0394	0.8401
Site	2	0.4920	2.7204	0.0694
Species	2	6.3539	35.135	0.0001
Sex	2	1.8742	10.364	0.0001
Year×Site	2	1.6892	9.3405	0.0001
Year×Species	2	0.4966	2.746	0.0654
Site×Species	4	0.8758	4.8426	0.0011
Year×Sex	1	0.0179	0.0988	0.7561
Site×Sex	4	0.5411	2.9922	0.0203
Species×Sex	4	0.2777	1.5357	0.1937
Year×Site×Species	4	2.1026	11.626	0.0001

<i>Year</i> × <i>Site</i> × <i>Sex</i>	2	0.0785	0.4341	0.6454
<i>Year</i> × <i>Species</i> × <i>Sex</i>	2	0.1651	0.9130	0.4087
<i>Site</i> × <i>Species</i> × <i>Sex</i>	8	0.8626	4.77	0.0001*
<i>Year</i> × <i>Site</i> × <i>Species</i> × <i>Sex</i>	4	0.0595	0.3289	0.8529
<i>Error</i>	351	0.1808		

236

237 **Animal movement**

238 The number of tagged animals that was detected after filtering and mean animal size for
 239 each year are shown in Table 3. In 2020, 21 animals tagged from the previous year were also
 240 detected (3 M lobsters; 3 F and 15 M crabs). In 2021, 10 animals tagged in 2020 were detected (1
 241 F and 1 M lobsters; 2 F and 6 M crabs). All animals tagged previously were included in the analyses
 242 for a given year.

243 **Table 3.** Total detections and mean size of American lobster (*Homarus americanus*) and rock crab
 244 (*Cancer irroratus*) at each year and tagging site after filtering.

	Species	Sex	Number of Individuals	Number of Detections	Size (mm) ± SE
Port-Mouton 2019	Lobster	M	19	39 000	86.24 ± 0.001
		F	10	23 682	83.32 ± 0.0004
	Rock crab	M	19	113 720	77.12 ± 0.003
		F	27	212 603	76.97 ± 0.008
Liverpool 2019	Lobster	M	24	99 990	88.03 ± 0.001
		F	19	78 448	88.86 ± 0.003
	Rock crab	M	33	269 358	109.49 ± 0.003
		F	12	118 642	86.99 ± 0.003
Liverpool 2020	Lobster	M	27	92 036	89.13 ± 0.002
		F	18	79 626	87.56 ± 0.008
	Rock crab	M	55	528 323	110.89 ± 0.002
		F	10	111 214	90.43 ± 0.002
Liverpool 2021	Lobster	M	21	80 611	89.60 ± 0.002
		F	21	76 597	86.98 ± 0.001
	Rock crab	M	27	260 245	94.37 ± 0.003
		F	24	184 193	88.10 ± 0.002


245

246

247 Movements of American lobster and rock crabs tagged in Liverpool Bay are highlighted in
248 Figure 3. There is a noted decrease in occupation of the farm site by lobsters over time whereas this
249 effect is not evident for rock crabs.

250 Residence time only varied as a function of “Species” ($p < 0.0001$; Table 4). Crabs stayed
251 longer in the acoustic array with a mean (\pm SE) residence time of 57.83 days \pm 2.41 compared to
252 31.80 days \pm 1.75 for lobsters (Figure 4) over the three-year study. For all years combined, a single
253 female crab tagged under the farm site in 2019 (100.81 mm CW) stayed the longest time within the
254 array (121.32 days). A male lobster tagged at Mersey Point in 2019 (86.4 mm size) remained the
255 shortest time (1.14 days).


256



257

258 **Figure 3.** Movement of American lobster (*Homarus americanus*) and rock crab (*Cancer*
 259 *irroratus*) detected within the acoustic array each year in Liverpool Bay, Nova Scotia. Each
 260 colour represents an individual animal.

261




262

263 **Figure 4.** Mean residence time (day \pm SE) by species for each tagging site and each year in
 264 Liverpool Bay, Nova Scotia. The colors represent the sex of the animals.

265 Distance travelled varied significantly as a function of “Year,” “Species,” and the “Site \times
 266 Species \times Sex” interaction (Table 4, Figure 5). In 2020, the mean (\pm SE) distance travelled by the
 267 two species was greater than that for the other years (47.18 ± 3.27 km compared to 35.94 ± 3.13
 268 km for 2019 and 33.69 ± 2.85 km for 2021) (Figure 5). Variation between the species is explained
 269 by the greater distance travelled by crabs (48.93 ± 2.77 km) relative to that by lobsters ($26.92 \pm$
 270 1.75) (Figure 5).


271 Variation in walking speed was only impacted by the factor “Species” (Table 4, Figure 6).
 272 Lobsters had a greater walking speed (1.75 ± 0.06 km/day) than crabs (1.22 ± 0.03 km/day).



273
 274 **Figure 5.** Mean distance travelled (\pm SE) by species from each tagging site and each year in
 275 Liverpool Bay, Nova Scotia. The colors represent the sex of the animals.

276

277



278

279 **Figure 6.** Mean walking speed (\pm SE) by species for each tagging site and each year in Liverpool
 280 Bay, Nova Scotia. The colors represent the sex of the animals.


281

282
 283 **Table 4.** Movement parameter results with the 4-way ANOVAs and PERMANOVA analyses for
 284 Liverpool Bay, Nova Scotia. Significant differences at $p < 0.05$ are indicated by bold font.

Source	ANOVA - Residence Time				ANOVA - Distance Travelled				PERMANOVA - Walking Speed			
	df	MS	F	p	df	MS	F	p	df	MS	F	p
Year	2	1592	2.322	0.1001	2	34.27	5.989	0.0029	2	0.0377	0.9863	0.3752
Site	2	62	0.091	0.9131	2	8.00	1.399	0.2487	2	0.0813	2.1272	0.1207
Species	1	49 664	72.430	1.4^{e-15}	1	184.60	32.259	3.6^{e-08}	1	1.0065	26.322	0.0001
Sex	1	312	0.455	0.5006	1	1.71	0.299	0.5847	1	0.0051	0.1342	0.7193
Year×Site	4	564	0.822	0.5119	4	2.59	0.453	0.7700	4	0.0055	0.1428	0.9662
Year×Species	2	470	0.686	0.5045	2	3.70	0.646	0.5250	2	0.0011	0.0298	0.9714
Site×Species	2	907	1.323	0.2682	2	4.22	0.737	0.4796	2	0.0054	0.1408	0.8622
Year×Sex	2	732	1.067	0.3455	2	6.44	1.125	0.3263	2	0.0070	0.1818	0.8302
Site×Sex	2	627	0.915	0.4019	2	9.12	1.593	0.2052	2	0.0130	0.3402	0.7146
Species×Sex	1	179	0.260	0.6103	1	3.24	0.566	0.4526	1	0.0079	0.2069	0.6514
Year×Site×Species	3	1096	1.598	0.1903	3	10.43	1.823	0.1434	3	0.0179	0.4682	0.6998
Year×Site×Sex	3	213	0.311	0.8177	3	1.35	0.237	0.8707	3	0.0320	0.8374	0.4837
Year×Species×Sex	2	1256	1.831	0.1623	2	4.51	0.787	0.4561	2	0.0264	0.6913	0.5015
Site×Species×Sex	2	2025	2.953	0.0539	2	17.43	3.045	0.0493	2	0.0123	0.3223	0.7208
Year×Site×Species×Sex	1	78	0.114	0.7362	1	2.42	0.423	0.5159	1	0.0292	0.7623	0.3834
Error	260	686			260	5.72			260	0.0382		

285
 286
 287 All movement parameters for the animals tagged in Port Mouton Bay varied as a function
 288 of tagging “Site” and “Species” (Table 5). Residence time for all the animals tagged in the north
 289 site differed significantly from that in the center and south and the time passed in the study area
 290 differed between lobsters and crabs with respectively 15.19 ± 2.86 days and 39.04 ± 4.30 days
 291 (Figure 7). The same pattern was observed for the distance travelled by animals tagged in the north
 292 site differing from that of animals tagged in the two other sites ($p = 0.0004$ for the center site and p
 293 < 0.0001 for the south site). The mean (\pm SE) distance travelled by the lobster differed from that
 294 for crabs (Figure 8). Walking speed differed between south and center sites ($p = 0.003$), and south

295 and north sites ($p < 0.0001$). Mean walking speed for lobsters was significantly different (Table 4,
296 Figure 8).



297
298 **Figure 7.** Mean movement parameters (\pm SE) by species for each tagging site in Port Mouton, Nova
299 Scotia. The colors represent the sex of the animals.


300 **Table 5.** Movement parameter results for the 3-way ANOVAs for Port Mouton, Nova Scotia.
301 Significant differences at $p < 0.05$ are indicated by bold font.

Source	Residence Time				Distance Travelled				Walking Speed			
	df	MS	F	p	df	MS	F	p	df	MS	F	p
Site	2	29.81	7.245	0.0015	2	31.443	14.138	8.47^{e-06}	2	1.386	11.008	7.94^{e-05}
Species	1	111.51	27.102	2.25^{e-06}	1	24.389	10.966	0.0015	1	9.162	72.750	4.25^{e-12}
Sex	1	5.37	1.305	0.25754	1	2.685	1.207	0.27602	1	0.035	0.281	0.598
Site×Species	2	1.23	0.299	0.74291	2	1.018	0.458	0.63469	2	0.143	1.136	0.328
Site×Sex	2	8.86	2.154	0.12444	2	2.899	1.303	0.27882	2	0.104	0.823	0.444
Species×Sex	1	1.73	0.420	0.51926	1	0.587	0.264	0.60914	1	0.049	0.387	0.536
Site×Species×Sex	2	9.46	2.299	0.10869	2	7.105	3.194	0.0477	2	0.148	1.174	0.316
Error	63	4.11			63	2.224			63	0.126		

302
303
304 **Farm attraction**


305 Variation between individual home ranges was significantly different as function of the Site
 306 × Species interaction such that the overlap with the farm area increases for rock crab over time
 307 whereas as that for lobster shows the opposite pattern (Table 6). The total home range for the three
 308 years for the lobsters tagged under the farm site differs from the crab home ranges (Figure 8).
 309 Overlap of crab and lobster distributions with the farm area are highlighted in Figure 9.

310 Home range for Port Mouton individuals varied as a function of “Site” and “Species” (Table
 311 7) such that there was a higher home range for lobster than rock crabs (Figure 10). For tagging sites,
 312 north and the south sites differed ($p = 0038$).



313
 314 **Figure 8.** Mean (\pm SE) home range by species for each tagging site and year in Liverpool Bay. The
 315 colors represent the sex of the animals.

316




317
 318 **Figure 9.** Mean (\pm SE) area overlap with fish farm by species for each tagging site and year in
 319 Liverpool Bay. The colors represent the sex of the animals.

320 **Table 6.** Results of PERMANOVAs for home range and the area that overlaps with the fish farm.
 321 Significant differences at $p < 0.05$ are indicated by bold font.

Source	Home range				Overlap with fish farm			
	df	MS	F	p	df	MS	F	p
Year	2	0.2037	0.4311	0.6508	2	0.6904	0.4966	0.6139
Site	2	0.0464	0.0981	0.9096	2	54.491	39.191	0.0001
Species	1	1.8079	3.826	0.0532	1	18.231	13.112	0.0003
Sex	1	0.0701	0.1484	0.6929	1	9.7329	7.0002	0.0097
Year×Site	4	0.1381	0.2922	0.8791	4	0.8343	0.6001	0.6639
Year×Species	2	0.0607	0.1284	0.8801	2	2.3159	1.6656	0.1919
Site×Species	2	1.8829	3.9848	0.0213	2	4.8358	3.478	0.0344
Year×Sex	2	0.0494	0.1044	0.9059	2	0.6401	0.4604	0.6393
Site×Sex	2	0.0690	0.1461	0.866	2	1.5641	1.125	0.3225
Species×Sex	1	0.5502	1.1644	0.2795	1	8.0102	5.7612	0.0189
Year×Site×Species	3	0.7779	1.6462	0.1848	3	3.1079	2.2353	0.0837
Year×Site×Sex	3	0.3236	0.6848	0.5642	3	3.4636	2.4911	0.063
Year×Species×Sex	2	0.0199	0.0422	0.9588	2	1.4864	1.0691	0.3469
Site×Species×Sex	2	0.6468	1.3688	0.2562	2	2.6662	1.9176	0.145

<i>Year</i> × <i>Site</i> × <i>Species</i> × <i>Sex</i>	1	0.4836	1.0233	0.3055	1	1.2318	0.8860	0.3417
<i>Error</i>	260	0.4725			260	1.3904		

322



323

324 **Figure 10.** Mean (\pm SE) home range by species for each tagging site in Port Mouton.

325 **Table 7.** Home range results for the 3-way ANOVAs for Port Mouton, Nova Scotia. Significant
 326 differences at $p < 0.05$ are indicated by bold font.

<i>Source</i>	<i>Home range</i>			
	<i>df</i>	<i>MS</i>	<i>F</i>	<i>p</i>
<i>Site</i>	2	388.7	5.658	0.0055
<i>Species</i>	1	2319.4	33.762	2.22^{e-07}
<i>Sex</i>	1	48.5	0.706	0.4041
<i>Site</i> × <i>Species</i>	2	70.2	1.022	0.3658
<i>Site</i> × <i>Sex</i>	2	8.7	0.126	0.8817
<i>Species</i> × <i>Sex</i>	1	9.3	0.135	0.7146
<i>Site</i> × <i>Species</i> × <i>Sex</i>	2	70.1	1.020	0.3664
<i>Error</i>	63	68.7		

327

328

329 **DISCUSSION**

330 This study evaluated the distribution (abundance and movement) of two decapods
331 (American lobster and rock crab) of commercial and ecological importance over a full production
332 cycle in Liverpool Bay and a few years post-salmonid production at a decommissioned finfish
333 aquaculture site in Port Mouton, Nova Scotia. In general terms, Atlantic salmon aquaculture in
334 Liverpool Bay was observed to affect the abundance and movement of both American lobster and
335 rock crabs.

336 Many organisms may be associated with finfish farms because of the physical structure they
337 provide and the trophic subsidy they offer in terms of lost feed and faeces (Callier et al. 2018).
338 While the bulk of this work has focused on fish, birds, and marine mammals (Barrett et al. 2019),
339 other taxonomic groups have also been the focus of studies, including benthic invertebrates (Callier
340 et al. 2018). Indeed, several studies have shown that decapods use and assimilate waste from marine
341 finfish farms. For example, northern shrimp (*Pandalus borealis*) fatty acid signatures were altered
342 close to farms relative to those caught distant from farms (Olsen et al. 2012). Likewise, caramote
343 prawn (*Melicertus kerathurus*) show isotopic evidence that animals close to the farm had been
344 feeding on farm waste (Izquierdo-Gomez et al. 2015). Woodcock et al. (2018) showed that brown
345 crab (*Cancer pagurus*) showed fatty acid and stable isotope evidence of being affected by the farm
346 at distances up to 1 km from the farm. A study on American lobster and rock crabs found that both
347 species had fatty acid profiles indicating that they had been feeding on feed waste or faeces by
348 comparing the results from animals in similar locations without fish farms (Sardenne et al. 2020).
349 Likewise, Baltadakis et al. (2020) showed that juvenile European lobster (*Homarus gammarus*) at
350 a control site differed those deployed adjacent to an Atlantic salmon farm in terms of fatty acid
351 signatures due to organic loading from the farm.

352 Several studies have evaluated the abundance of decapods around fish farms in eastern
353 Canada with a focus on lobster. For example, Lawton (2002) observed that lobster in the area of a
354 Grand Manan, New Brunswick, Canada, salmon farm were less abundant during a period when
355 farms were operational, particularly for egg-bearing (berried) lobster, than in periods when salmon
356 were not in fish cages and that historical patterns of site occupation returned when the farm was
357 removed. Milewski et al. (2018) suggested that lobster catches of both commercial and berried
358 females decreased during farming operations at a Port Mouton, Nova Scotia, Canada salmonid
359 farm. Likewise, Wiber et al. (2012) suggest that fishers in the Bay of Fundy, New Brunswick,
360 believe that berried female lobster avoid areas where salmon aquaculture has established. However,
361 Grant et al. (2016) suggested that this is not the case and that longer-term studies suggest that
362 salmon farms have no obvious impact on lobster abundance. For example, Grant et al. (2019)
363 sampled lobster under a Grand Manan salmon farm over 8 years and at appropriate reference areas
364 and did not observe variation in lobster abundances between the two habitat types (farm and
365 reference) for either market size or berried females. The present study noted a marked decrease in
366 the abundance of lobster from the fallow year relative to the year when the farm contained two-
367 year-old fish, although this trend was not observed in reference areas (Fralick Cove and Mersey
368 Point). In contrast, this effect was not observed for rock crabs. Milewski et al. (2018) suggested
369 that the salmonid farm in Port Mouton created benthic conditions due to excess feed that were
370 unfavorable to lobsters. However, Milewski et al. (2018) invoke mechanisms occurring at a larger
371 spatial scale (e.g. hypoxia and sulphide levels, as outlined in Hargrave et al. 1997, for example)
372 than patterns observed in the present study and the data to support predicted effects in Milewski et
373 al. (2018) show the opposite pattern than would be predicted (i.e. there were fewer berried lobster
374 in the region surrounding the farm in fallow years than in years when fish were in cages).

375 Lobster and crab movements differed and varied spatially across years in Liverpool Bay and
376 spatially in Port Mouton. As for lobster within mussel leases (Lavoie et al. 2022, Lees et al. 2023),
377 lobster in Liverpool Bay showed little affinity to the salmon farm there as the lobster caught and
378 released adjacent to or below the farm did not stay in the area in the fallow year and this effect only
379 increased in subsequent production years. This is reflected by the spatial overlap of lobster
380 distribution with the salmon farm, which declined over time. In contrast, rock crabs seem to be
381 associated with farms, even in the fallow year. This is likely due to fall-off of fouling organisms
382 (mussels *Mytilus edulis*) in the fallow year (McKindsey, personal observations) and consuming
383 salmon feed during production years (Sardenne et al. 2020). The effect of mussel fall-off on rock
384 crab distributions was previously noted in Prince Edward Island by Lees et al. (2023) and rock crabs
385 are known to consume salmon feed from laboratory studies (Drolet et al. 2022). Thus, as opposed
386 to lobster, the spatial overlap of crab distributions increased over time.

387 Other movement metrics also varied by species. Both walking speed and home range in
388 Liverpool Bay were greater for lobster than for rock crab, as was also observed for these species in
389 the areas surrounding mussel culture sites in Prince Edward Island (Lees et al. 2023). The same
390 effect was also observed in Port Mouton. Other movement metrics (distance travelled and residence
391 time) were both lower for lobster than for rock crab as these are reflected by the quicker movement
392 of lobsters leaving the acoustic arrays established in both study sites more rapidly than rock crabs.

393 Given the results from this study, it is unclear what effects salmonid aquaculture may have
394 on decapods in the surrounding area. While both American lobster and rock crab uptake nutrients
395 from aquaculture activities (Sardenne et al. 2020), it is also known that a diet of only salmon feed
396 may have negative impacts on rock crab condition (Drolet et al. 2022). However, this clearly does
397 not occur under field conditions as crabs were clearly attracted to fallen mussels in the farm area in

398 the fallow year. Similar work on lobster is underway (Drolet, personal communications). Variation
399 in the overlap of lobster distributions with the farm area over time indicates that lobsters are less
400 abundant in the farm area as aquaculture production increases. However, there was no indication
401 that movement of lobsters in areas adjacent to the farm diminished. Likewise, the spatial overlap of
402 rock crabs with the farm increased over time, although the movement of animals adjacent to the
403 farm remained similar over the three years of the study. As for Bay of Fundy aquaculture sites
404 (Walters 2007), salmonid aquaculture will continue to interact with decapods throughout eastern
405 Canada. It is hoped that the results presented here will inform the discussion on the importance of
406 such interactions.

407

408 **ACKNOWLEDGEMENTS**

409 We are grateful to Émilie Simard, Frédéric Hartog, Jean-Daniel Tourangeau-Larivière,
410 Simon Jacques, and Delphine Cottier from Fisheries and Oceans Canada, and Todd Mosher, David
411 Cook, and Jennifer Feehan from the Nova Scotia Department of Fisheries and Aquaculture for their
412 help in project planning and field assistance. This study was supported by Fisheries and Oceans
413 Canada and the Nova Scotia Department of Fisheries and Aquaculture. We also thank the fish
414 farmer collaborator, Cooke Aquaculture, for allowing access to their site.

415

416 **REFERENCES CITED**

- 417 Anderson MJ (2001) A new method for non-parametric multivariate analysis of variance. *Austral*
418 *Ecol* 26:32-45 doi: 10.1111/j.1442-9993.2001.01070.pp.x
- 419 Baltadakis A, Casserly J, Falconer L, Sprague M, Telfer TC (2020) European lobsters utilise
420 Atlantic salmon wastes in coastal integrated multi-trophic aquaculture systems. *Aquacult*
421 *Environ Interact* 12:485-494 doi: 10.3354/aei00378
- 422 Barrett LT, Swearer SE, Dempster T (2019) Impacts of marine and freshwater aquaculture on
423 wildlife: a global meta-analysis. *Rev Aquacult* 11:1022-1044 doi: 10.1111/raq.12277
- 424 Calenge, C. (2006) The package adehabitat for the R software: a tool for the analysis of space and
425 habitat use by animals. *Ecol Modell* 197:516-519 doi: 10.1016/j.ecolmodel.2006.03.017
- 426 Callier MD, Byron CJ, Bengtson DA, Cranford PJ, Cross ST, Focken U, Jansen HM, Kamermans
427 P, Kiessling A, Landry T, O'Beirn F, Petersson E, Rheault RB, Strand Ø, Sundell K, Svåsand
428 T, Wikfors GH, McKindsey CW (2018) Attraction and repulsion of mobile wild organisms to
429 finfish and shellfish aquaculture: a review. *Rev Aquacult* 10:924-949 doi: 10.1111/raq.12208
- 430 Drolet D, Riley C, Robert S, Estrada R, Gianasi BL, McKindsey CW (2022) Effect of aquaculture-
431 related diets on the long-term performance and condition of the rock crab, *Cancer irroratus*.
432 *Front Mar Sci* 9:865390 doi: 10.3389/fmars.2022.865390
- 433 Espinoza M, Farrugia TJ, Webber DM, Smith F, Lowe CG (2011) Testing a new acoustic telemetry
434 technique to quantify long-term, fine-scale movements of aquatic animals. *Fish Res*
435 108:364–371 doi: 10.1016/j.fishres.2011.01.011

- 436 FAO Fisheries and Aquaculture Department (2022) The state of world fisheries and aquaculture
437 2022. Towards Blue Transformation. Food and Agriculture Organization of the United Nations,
438 Rome, xxv + 236 p.
- 439 Fisheries and Oceans Canada (2021) Aquaculture Production and Value. [https://www.dfo-](https://www.dfo-mpo.gc.ca/stats/aqua/aqua21-eng.htm)
440 [mpo.gc.ca/stats/aqua/aqua21-eng.htm](https://www.dfo-mpo.gc.ca/stats/aqua/aqua21-eng.htm) (accessed October 10, 2023)
- 441 Fisheries and Oceans Canada (2022) Atlantic rock crab. [https://www.dfo-mpo.gc.ca/species-](https://www.dfo-mpo.gc.ca/species-especies/profiles-profil/atl-rock-crab-crabe-commun-eng.html)
442 [especies/profiles-profil/atl-rock-crab-crabe-commun-eng.html](https://www.dfo-mpo.gc.ca/species-especies/profiles-profil/atl-rock-crab-crabe-commun-eng.html) (accessed January 18, 2024)
- 443 Florko KRN, Davidson E, R., Lees KJ, Hammer LJ, Lavoie M-F, Lennox R, Simard É,
444 Archambault P, Auger-Méthé M, McKindsey CW, Whoriskey FG, Furey NB (2021) Tracking
445 movements of decapod crustaceans: a review of a half-century of telemetry-based studies. *Mar*
446 *Ecol Prog Ser* 679:219-239 doi: 10.3354/meps13904
- 447 Gendron L, Fradette P, Godbout G (2001) The importance of rock crab (*Cancer irroratus*) for
448 growth, condition and ovary development of adult American lobster (*Homarus americanus*). *J*
449 *Exp Mar Biol Ecol* 262:221-241 doi: 10.1016/S0022-0981(01)00297-0
- 450 Grant J, Filgueira R, Barell J (2016) Lack of interaction between finfish aquaculture and lobster
451 catch in coastal Nova Scotia. *Mar Poll Bull* 110:613-615 doi: 10.1016/j.marpolbul.2016.06.043
- 452 Grant J, Simone M, Daggett T (2019) Long-term studies of lobster abundance at a salmon
453 aquaculture site, eastern Canada. *Can J Fish Aquat Sci* 76:1096-1102 doi: 10.1139/cjfas-2017-
454 0547
- 455 Hanson JM (2009) Predator-prey interactions of American lobster (*Homarus americanus*) in the
456 southern Gulf of St. Lawrence, Canada. *N Z J Mar Freshwater Res* 43:69-88 doi:
457 10.1080/00288330909509983

- 458 Hargrave BT, Phillips GA, Doucette LI, White MJ, Milligan TG, Wildish DJ, Cranston RE (1997)
459 Assessing benthic impacts of organic enrichment from marine aquaculture. *Water, Air, Soil*
460 *Pollut* 99:641-650 doi: 10.1007/BF02406903
- 461 Horricks RA, Lewis-McCrea LM, Reid GK (2022) Interactions between American lobster
462 (*Homarus americanus*) and salmonid aquaculture in the Canadian Maritimes. *Can J Fish Aquat*
463 *Sci* 79:1561–1571 doi: 10.1139/cjfas-2021-0252
- 464 Izquierdo-Gomez D, Sanchez-Jerez P, Bayle-Sempere JT, Loader NJ, Garcia de Leaniz C (2015)
465 Effects of coastal fish farms on body size and isotope composition of wild penaeid prawn. *Fish*
466 *Res* 172:50-56 doi: 10.1016/j.fishres.2015.06.017
- 467 Lavoie MF, Simard É, Drouin A, Archambault P, Comeau LA, Mckindsey CW (2022) Movement
468 of American lobster *Homarus americanus* associated with offshore mussel *Mytilus edulis*
469 aquaculture. *Aquacult Environ Interact* 14:189-204 doi: 10.3354/aei00437
- 470 Lawton P (2002) Prior evaluation of sensitive lobster fishery habitat in relation to salmon
471 aquaculture and new monitoring approaches. In: Hargrave BT (ed) *Environmental Studies for*
472 *Sustainable Aquaculture (ESSA): 2002 Workshop Report Can Tech Rep Fish Aquat Sci #2411,*
473 *p 10-15*
- 474 Lees KJ, Lavoie M-F, MacGregor KA, Simard É, Drouin A, Comeau LA, McKindsey CW (2023)
475 Movement of American lobster *Homarus americanus* and rock crabs *Cancer irroratus* around
476 mussel farms in Malpeque Bay, Prince Edward Island, Canada. *Aquacult Environ Interact*
477 15:179–193doi: 10.3354/aei00461
- 478 Lees KJ, Mill AC, Skerritt DJ, Robertson PA, Fitzsimmons C (2020) Spatial overlap, proximity,
479 and interaction between lobsters revealed using acoustic telemetry. *Mar Ecol Prog Ser* 645:109-
480 124 Doi: 10.3354/meps13376

- 481 Loucks RH, Smith RE, Fisher EB (2014) Interactions between finfish aquaculture and lobster
482 catches in a sheltered bay. *Mar Poll Bull* 88:255-259 doi: 10.1016/j.marpolbul.2014.08.035
- 483 Loucks RH, Smith RE, Fisher EB (2016) A response to the letter to the editor “Lack of interaction
484 between finfish aquaculture and lobster catch in coastal Nova Scotia”. *Mar Poll Bull* 110:616-
485 618 doi: 10.1016/j.marpolbul.2016.06.043
- 486 McKee A, Grant J, Barrell J (2021) Mapping American lobster (*Homarus americanus*) habitat for
487 use in marine spatial planning. *Can J Fish Aquat Sci* 78:704-720 doi: 10.1139/cjfas-2020-0051
- 488 Milewski I, Loucks RH, Fisher B, Smith RE, McCain JSP, Lotze HK (2018) Sea-cage aquaculture
489 impacts market and berried lobster (*Homarus americanus*) catches. *Mar Ecol Prog Ser* 598:85-
490 97 doi: 10.3354/meps12623
- 491 Olsen SA, Ervik A, Grahl-Nielsen O (2012) Tracing fish farm waste in the northern shrimp
492 *Pandalus borealis* (Krøyer, 1838) using lipid biomarkers. *Aquacult Environ Interact* 2:133-144
493 doi: 10.3354/aei00036
- 494 Orrell DL, Hussey NE (2022) Using the VEMCO Positioning System (VPS) to explore fine-scale
495 movements of aquatic species: applications, analytical approaches and future directions. *Mar*
496 *Ecol Prog Ser* 687:195-216 doi: 10.3354/meps14003
- 497 Piper DJW, Mudie PJ, Letson JRJ, Barnes NE, Iuliucci RJ (1986) 1986. The marine geology of the
498 inner Scotian shelf off the South Shore, Nova Scotia. Geological Survey of Canada, Ottawa
- 499 Quinn G, Keough M (2002) Experimental design and data analysis for biologists. Cambridge
500 University Press, Cambridge
- 501 R Core Team (2023) R: A language and environment for statistical computing. R Foundation for
502 Statistical Computing, Vienna, Austria, <https://www.R-project.org/>

- 503 Sardenne F, Simard M, Robinson SMC, McKindsey CW (2020) Consumption of organic wastes
504 from coastal salmon aquaculture by wild decapods. *Sci Total Environ* 711:134863 doi:
505 10.1016/j.scitotenv.2019.134863
- 506 Skerritt DJ, Robertson PA, Mill AC, Polunin NVC (2015) Fine-scale movement, activity patterns
507 and home-ranges of European lobster *Homarus gammarus*. *Mar Ecol Prog Ser* 536:203-219
508 doi: 10.3354/meps11374
- 509 Signer J, Fieberg J, Avgar T (2019) Animal movement tools (amt): R package for managing
510 tracking data and conducting habitat selection analyses. *Ecol Evol* 9:880-890 doi:
511 10.1002/ece3.4823
- 512 Walters BB (2007) Competing use of marine space in a modernizing fishery: salmon farming meets
513 lobster fishing on the Bay of Fundy. *Can Geogr* 51:139-159 doi: 10.1111/j.1541-
514 0064.2007.00171.x
- 515 Wiber MG, Young S, Wilson L (2012) Impact of aquaculture on commercial fisheries: Fishermen's
516 local ecological knowledge. *Hum Ecol*, doi: 10.1007/s10745-011-9450-7

Supplementary Material

3.1.1 2019 – Liverpool and Port-Mouton




Figure X. Tracks for animals tagged in the North site, Port-Mouton 2019. The black dots indicate receiver locations.




Figure X. Tracks for animals tagged in the Center site, Port-Mouton 2019. The black dots indicate receiver locations.




Figure X. Tracks for animals tagged in the South site, Port-Mouton 2019. The black dots indicate receiver locations.




Figure X. Tracks for animals tagged at Mersey Point, Liverpool 2019. The black dots indicate receivers and the farm is represented by the black polygon.




Figure X. Tracks for animals tagged at Fralick Cove, Liverpool 2019. The black dots indicate receivers and the farm is represented by the black polygon.




Figure X. Tracks for lobsters tagged near the fish farm area, Liverpool 2019. The black dots indicate receivers and the farm is represented by the black polygon.




Figure X. Tracks for crabs tagged near the fish farm area, Liverpool 2019. The black dots indicate receivers and the farm is represented by the black polygon.

3.1.2 2020 – Liverpool




Figure X. Tracks for animals tagged at Mersey Point, Liverpool 2020. The black dots indicate receivers and the farm is represented by the black polygon.




Figure X. Tracks for animals tagged at Fralick Cove, Liverpool 2020. The black dots indicate receivers and the farm is represented by the black polygon.




Figure X. Tracks for lobsters tagged near the fish farm area, Liverpool 2020. The black dots indicate receivers and the farm is represented by the black polygon.




Figure X. Tracks for crabs tagged near the fish farm area, Liverpool 2020. The black dots indicate receivers and the farm is represented by the black polygon.

3.1.3 2021 – Liverpool




Figure X. Tracks for animals tagged at Mersey Point, Liverpool 2021. The black dots indicate receivers and the farm is represented by the black polygon.




Figure X. Tracks for animals tagged at Fralick Cove, Liverpool 2021. The black dots indicate receivers and the farm is represented by the black polygon.




Figure X. Tracks for lobsters tagged near the fish farm area, Liverpool 2021. The black dots indicate receivers and the farm is represented by the black polygon.




Figure X. Tracks for crabs tagged near the fish farm area, Liverpool 2021. The black dots indicate receivers and the farm is represented by the black polygon.